



Scientific note

**Predation of *Ameivula ocellifera* (Squamata, Teiidae) by *Oxybelis aeneus* (Squamata, Colubridae), in the Caatinga, Northeastern Brazil, including a list of saurophagy by this snake**

Natallia Vívian N. da Silva<sup>1,\*</sup>, Anna Virginia A. de Mello<sup>1</sup>, Patricia Marques do A. Oliveira<sup>1,2</sup>, José Henrique de Andrade Lima<sup>1,2</sup>, Lucas Rodrigues de Castro Ribeiro<sup>1</sup> & Pedro Ivo Simões<sup>1,2</sup>.

The most common method of identifying the feeding habits of a species is the dissection of the digestive tracts of animals that have been deposited in collections and/or museums or through occasional encounters (e. g. Prudente *et al.* 1998, Bernarde *et al.* 2000, Bovo & Sueiro 2012, Dorigo *et al.* 2014). Encounters with snakes in their natural environments are rare, and the observation of feeding events is especially uncommon (Mushinsky 1987). However, the recording of this kind of ecological interaction in nature is essential for improving our understanding of predator-prey relationships (Sousa *et al.* 2020). Herein we present a predation event of *Oxybelis aeneus* (Wagler, 1824) on *Ameivula ocellifera* (Spix, 1825) in a Caatinga area in northeastern Brazil.

*Oxybelis aeneus* is a diurnal arboreal, midsize colubrid snake with opisthognath dentition (Savage 2002) and demonstrates "sit-and-wait" foraging behavior, as well as oviparous reproduction (Henderson and Nickerson 1976; Franzini *et al.* 2018). This species is distributed throughout southern North America, Central America, and in

most of the South American continent (Savage 2002, Uetz 2017), and is commonly found in the semi-arid Caatinga biome in northeastern Brazil (Vanzolini *et al.* 1980, Guedes *et al.* 2014, Costa *et al.* 2018). It is easily recognized by its slim body, elongated snout, and gray and brown dorsum, without lateral stripes (Keiser 1974). This combination of features allows individuals to camouflage themselves among dry branches (Mesquita *et al.* 2012). Its diet consists of small vertebrates such as lizards, amphibians, birds, small mammals, and fish (Vanzolini *et al.* 1980, Hetherington 2006, Grant & Lewis 2010, Santos *et al.* 2012, Oliveira *et al.* 2020).

*Ameivula ocellifera* is a small-medium diurnal lizard (Vitt & Caldwell 2009) with active ground foraging, characterized by constant movement and exploratory behavior (Pianka 1966), and is more active during the hottest hours of the day (Sales & Freire 2015). Its body color varies from green to light brown with longitudinal dotted lines (Lisboa *et al.* 2017). *A. ocellifera* is widely distributed throughout the Caatinga, Cerrado and Restinga habitats along the northeastern coast of Brazil (Mesquita & Colli 2003).

<sup>1</sup> Centro de Biociências, Laboratório de Herpetologia, Departamento de Zoologia, Universidade Federal de Pernambuco, 50670-901 Recife, Pernambuco, Brazil.

<sup>2</sup> Programa de Pós-Graduação em Biologia Animal, Centro de Biociências, Universidade Federal de Pernambuco, 50670-901 Recife, Pernambuco, Brazil.

\*Corresponding author: [natallivivian8@gmail.com](mailto:natallivivian8@gmail.com)

The observation occurred on 12th November 2020, at 10:47 h, in the Parque Nacional do Catimbau (-8.575194 S, -37.246083 W; 756 m), a conservation unit in the Caatinga, located in the municipality of Buíque, State of Pernambuco, northeastern Brazil. An *O. aeneus* individual (SVL 90 cm) was sighted, with the posterior half of its body supported on the thorny branches of a shrub (~1.50 m above ground) and the anterior half suspended perpendicular to the ground, with an adult *A. ocellifera* female (SVL ~6.5 cm) immobilized in its mouth, approximately 1.30 m above ground. The lizard was held by its head, up to its jaw articulation (Figure 1A). After 12 min, the snake lifted its prey up to the height of the branches, approximately 1.40 m above ground (Figure 1B). After another 09 min, the snake moved the lizard 10 cm below the branch where it was situated and began to ingest its prey (Figure 1C). After 02 min, the snake took its prey to the branches again and ingested it completely.

Among the 14 lizard species recorded as *O. aeneus* prey, at least six are sit-and-wait foragers, four are active foragers and one is intermediate forager (Table 1). According to Sellmeijer & Van den Burg. (2020), the diversity of foraging classifications of *O. aeneus* prey reinforces its generalist feeding behavior. Additionally, nine species are terrestrial (Table 1), which demonstrates that although *O. aeneus* is an arboreal snake, it efficiently preys upon lizards on the ground.

Although saurophagy by colubrid snakes is well-reported in the literature (e.g., Vanzolini *et al.* 1980, Vitt & Vangilder 1983, Mikalauskas *et al.* 2017, Lissa *et al.* 2018), only a few studies describe the behavior of these snakes during predation events in detail (Almeida *et al.* 2009, Franzini *et al.* 2018, Santana & Teixeira 2020, Sousa *et al.* 2020), which



**Figure 1.** *Oxybelis aeneus* preying on an *Ameivula ocellifera* adult at the Parque Nacional do Catimbau, state of Pernambuco, northeastern Brazil, on 12th November 2020. Chronology of the field observation: (A) at the moment of the first sighting of the snake; (B) 12 minutes after encounter, and (C) 21 minutes after encounter.

can be exemplified by the available literature on *O. aeneus* diet (Table 1). Despite *A. ocellifera* has already been registered as a prey item of *O. aeneus* (Mesquita *et al.* 2012), no behavioral observation has been provided. In the present record, we observed behavioral steps common to other *O. aeneus* predation events on other lizards. The snake was

positioned on top of branches, with its body partly positioned perpendicular to the ground, and it immobilized and ingested its prey starting at the head (Almeida *et al.* 2009, Santana & Teixeira 2020, Sousa *et al.* 2020). Thus, our report suggests a stereotyped predation strategy among different populations of this species.

**Table 1.** Lizard species that *Oxybelis aeneus* preys upon, with its respective foraging modes, habits, and references. \*Papers with detailed predation behavior.

Family	Prey	Foraging	Habit	Reference
<b>Corytophanidae</b>	<i>Basiliscus vittatus</i>	Active	Arboreal	Henderson 1982
<b>Dactyloidae</b>	<i>Anolis</i> sp.	-	Arboreal	Henderson 1982
<b>Gekkonidae</b>	<i>Hemidactylus mabouia</i>	Sit-and-Wait	Arboreal	Mesquita <i>et al.</i> 2012; Franzini <i>et al.</i> 2018*
<b>Phyllodactylidae</b>	<i>Gymnodactylus geckoides</i>	Sit-and-Wait	Terrestrial	Oliveira <i>et al.</i> 2020
<b>Polychrotidae</b>	<i>Polychrus acutirostris</i>	Sit-and-Wait	Arboreal	Oliveira <i>et al.</i> 2020
<b>Mabuyidae</b>	<i>Brasilicincus heathi</i>	Intermediate	Terrestrial	Mesquita <i>et al.</i> 2012; Oliveira <i>et al.</i> 2020
	<i>Copeoglossus arajara</i>	-	Terrestrial	Oliveira <i>et al.</i> 2020
<b>Sphaerodactylidae</b>	<i>Coleodactylus meridionalis</i>	-	Terrestrial	Oliveira <i>et al.</i> 2020
<b>Teiidae</b>	<i>Ameivula ocellifera</i>	Active	Terrestrial	Mesquita <i>et al.</i> 2012; This paper*
	<i>Ameivula pyrrhogularis</i>	Active	Terrestrial	Oliveira <i>et al.</i> 2020
	<i>Aspidoscelis angusticeps</i>	Active	Terrestrial	Henderson 1982
<b>Tropiduridae</b>	<i>Tropidurus cocorobensis</i>	Sit-and-Wait	Terrestrial	Almeida <i>et al.</i> 2009*
	<i>Tropidurus hygomi</i>	Sit-and-wait	Terrestrial	Santos <i>et al.</i> 2012
	<i>Tropidurus hispidus</i>	Sit-and-Wait	Semi-arboreal	Mesquita <i>et al.</i> 2012; Sousa <i>et al.</i> 2020*; Santana & Teixeira 2020*; Oliveira <i>et al.</i> 2020

## ACKNOWLEDGEMENTS

We are grateful for PELD Catimbau-CNPq (Long Term Ecological Project - Catimbau), and for research scholarship from Pró-Reitoria de Pesquisa e Inovação of UFPE (PROPesqi) and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq, Edital Propesqi nº 08/2020 PIBIC-AF/UFPE/CNPQ) to NVNS. Additional scholarships are provided by Fundação de

Amparo à Ciência e Tecnologia do Estado de Pernambuco (FACEPE, IBPG - 1566 - 2.04/19) to PMAO and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (Capes, 88887.511079/2020-00) to JHAL.

## REFERENCES

Almeida, G.V., Silva, G.L., Campos, T.F., Muniz, S.L. & Santos, E.M. (2009) Predação do lagarto *Tropidurus*

- cocorobensis* pela serpente *Oxybelis aeneus*. *Boletim do Museu de Biologia Mello Leitão* (N.Sér.) 25: 83–86.
- Bernarde, P.S., Moura-Leite, J.C., Machado, R.A. & Kokubum, M.N.C. (2000) Diet of the colubrid snake, *Thamnodynastes strigatus* (Günther, 1858) from Paraná State, Brazil, with field notes on anuran predation. *Revista Brasileira de Biologia* 60(4): 695–699. <https://doi.org/10.1590/s0034-7108200000400022>
- Bovo, R.P. & Sueiro, L.R. (2012) Records of predation on *Itapotihyla langsdorffii* (Anura: Hylidae) by *Chironius bicarinatus* (Serpentes: Colubridae) with notes on foraging substrate. *Herpetology Notes* 5: 291–292.
- Costa, T.B., Laranjeiras, D.O., Caldas, F.L.S., Santana, D.O., Silva, C.F., Alcântara, E.P., Brito, S.V., Galdino, J.Y., Mesquita, D.O., Faria, R.G., França, F.G.R., Ávila, R.W. & Garda, A.A. (2018) Herpetofauna of protected areas in the Caatinga VII: Aiuaba Ecological Station (Ceará, Brazil). *Herpetology Notes* 11: 929–941.
- Dorigo, T.A., Vrcibradic, D., Borges-Junior, V.N.T. & Rocha, C.B.D. (2014) New records of anuran predation by snakes of the genus *Thamnodynastes* Wagler, 1830 (Colubridae: Dipsadinae) in the Atlantic rainforest of southeastern Brazil. *Herpetology Notes* 7: 261–264.
- Franzini, L.D., Pedro, C.K.B., Queiroga Cavalcanti, L.B. & Mesquita, D.O. (2018) Predation of *Hemidactylus mabouia* (Sauria: Gekkonidae) by a vine snake *Oxybelis aeneus* (Serpentes: Colubridae) in an Atlantic Forest fragment, northeastern Brazil. *Pesquisa e Ensino em Ciências Exatas e da Natureza* 2(1): 67–70. <https://doi.org/10.29215/pecen.v2i1.587>
- Grant, P.B. & Lewis, T.R. (2010) Predation attempt by *Oxybelis aeneus* (Wagler) (Mexican Vine-snake) on *Basiliscus plumifrons* (Cope). doi: [https://doi.org/10.13128/Acta\\_Herpetol-8531](https://doi.org/10.13128/Acta_Herpetol-8531)
- Guedes, T.B., Nogueira, C. & Marques, O.A.V. (2014) Diversity, natural history, and geographic distribution of snakes in the Caatinga, Northeastern Brazil. *Zootaxa* 3863(1): 1–93. <https://doi.org/10.11646/zootaxa.3863.1.1>
- Henderson R.W. and M.A. Nickerson (1976). Observations on the behavioral ecology of three species of Imantodes (Serpentes: Colubridae). *Journal of Herpetology* 10: 205–210. <https://doi.org/10.2307/1562981>
- Henderson, R.W. (1982) Trophic relationships and foraging strategies of some new world tree snakes (*Leptophis*, *Oxybelis*, *Uromacer*). *Amphibia-Reptilia* 3: 71–80. <https://doi.org/10.1163/156853882x00185>
- Hetherington, T.E. (2006) *Oxybelis aeneus* diet. *Herpetological Review* 37: 94–95.
- Keiser Jr., E.D. (1974) A systematic study of the Neotropical Vine Snake *Oxybelis aeneus*. *Bulletin of Texas Memorial Museum* 22: 1–51.
- Lisboa, C. M., Bajer, K., Pessoa, D. M., Huber, M. A. & Costa, G. C. (2017) Female Brazilian whiptail lizards (*Cnemidophorus ocellifer*) prefer males with high ultraviolet ornament reflectance. *Behavioural processes* 142: 33–39. <https://doi.org/10.1016/j.beproc.2017.05.009>
- Lissa, D.F., Carmem, K.B.P., Lucas, B.Q.C. & Daniel, O.M. (2018) Predation of *Hemidactylus mabouia* (Sauria: Gekkonidae) by a vine snake *Oxybelis aeneus* (Serpentes: Colubridae) in an Atlantic Forest fragment, Northeastern Brazil. *Pesquisa e Ensino em Ciências Exatas e da Natureza* 2: 67–70. <https://doi.org/10.29215/pecen.v2i1.587>
- Mesquita, D.O. & Colli, G.R. (2003) The ecology of *Cnemidophorus ocellifer* (Squamata, Teiidae) in a neotropical savanna. *Journal of Herpetology* 37(3): 498–509. <https://doi.org/10.1670/179-02a>
- Mesquita, P. C., Borges-Nojosa, D. M., Passos, D. C. & Bezerra, C. H. (2012) Activity patterns of the Brown Vine snake *Oxybelis aeneus* (Wagler, 1824) (Serpentes, Colubridae) in the Brazilian semi-arid. *Animal Biology* 62(3): 289–299. <https://doi.org/10.1163/157075611x618228>
- Mikalauskas, J. S., Santana, D. O., & Ferrari, S. F. (2017). Lizard predation *Tropidurus hispidus* (Squamata, Tropiduridae) by false coral snake *Oxyrhopus trigeminus* (Squamata, Dipsadidae) in the Caatinga, in northeastern Brazil. *Pesquisa e Ensino em Ciências Exatas e da Natureza*, 1(1). doi: <http://dx.doi.org/10.29215/pecen.v1i1.174>
- Mushinsky, H.R. (1987) Foraging ecology. In: Seigel, R.A., Collins, J.T. & Novak, S.S. (Eds) *Snakes: ecology and*

- evolutionary biology*. MacMillan Public, New York, pp. 302–334. <https://doi.org/10.2307/1445695>
- Oliveira, C.R., Brasileiro, A.C. & Mascarenhas, W. (2020) *Oxybelis aeneus* (Brown Vine Snake). Diet *Herpetological review* 51(4): 876.
- Pianka, E.R. (1966) Convexity, desert lizards, and spatial heterogeneity. *Ecology* 47: 1055–1059.
- Prudente, A.L.D.C., Moura-Leite, J.C.D. & Morato, S.A.A. (1998) Alimentação das espécies de *Siphlophis Fitzinger* (Serpentes, Colubridae, Xenodontinae, Pseudoboini). *Revista Brasileira de Zoologia* 15(2): 375–383. <https://doi.org/10.2307/1935656>
- Sales, R.F.D. & Freire, E.M.X. (2015) Diet and foraging behavior of *Ameivula ocellifera* (Squamata: Teiidae) in the Brazilian semiarid Caatinga. *Journal of Herpetology* 49(4): 579–585. <https://doi.org/10.1670/14-041>
- Santana, D.O., & Teixeira, A.A.M. (2020) Predation of the lizard *Tropidurus hispidus* (Squamata, Tropiduridae) by the vine snake *Oxybelis aeneus* (Serpentes: Colubridae) in the Caatinga, northeastern Brazil. *Pesquisa e Ensino em Ciências Exatas e da Natureza* 4: 01–06. <https://doi.org/10.29215/pecen.v4i0.1327>
- Santos, R.A., Santana, D.O., Caldas, F.L.S. & Faria, R.G. (2012) *Tropidurus hygomi* (Reinhardt's lava lizard) Predation. *Herpetological Review* 43: 490–491.
- Savage, J.M. (2002) *The amphibians and reptiles of Costa Rica: a herpetofauna between two continents, between two seas*. University of Chicago Press, Chicago, 954 pp.
- Sellmeijer, B. & van den Burg, M. P. (2020) Tadpole predation in the chemically defended *Oophaga pumilio* (Anura: Dendrobatidae) by *Oxybelis aeneus* (Squamata: Colubridae). *Herpetology Notes* 13: 301–303.
- Sousa, J.D., Andrade Lima, J.H. & Kokubum, M.N. (2020) Predation of *Tropidurus hispidus* (Squamata, Tropiduridae) by *Oxybelis aeneus* (Squamata, Colubridae) in two biomes in the state of Pernambuco, Northeast of Brazil. *Herpetology Notes* 13: 171–173.
- Uetz P. (2017) *Oxybelis aeneus*. *The Reptile Database*. Available at: <http://www.reptiledatabase.org> (Accessed on: 04 June 2021).
- Vanzolini, P.E., Ramos-Costa, A.M.M. & Vitt, L.J. (1980) *Répteis das caatingas*. Academia Brasileira de Ciências, Rio de Janeiro, 161 pp. <https://doi.org/10.5962/bhl.title.109659>
- Vitt, L.J. & Caldwell, J.P. (2009) *Herpetology: an introductory biology of amphibians and reptiles*. Academic Press, New York, 720 pp.
- Vitt, L.J. & Vangilder, L.D. (1983) Ecology of a snake community in northeastern Brazil. *Amphibia-Reptilia* 4: 273–296. <https://doi.org/10.1163/156853883X00148>