









Scientific note

**New records of *Boana raniceps* (Cope, 1862) (Anura, Hylidae) in Distrito Federal, Central Brazil: evidence of recent distribution expansion?**

Beatriz Diogo Vasconcelos<sup>1,2\*</sup> , Emayre Rebecca Campos<sup>1,3</sup> , Vitor Matheus Alcântara de Sena<sup>1,4</sup> ,  
Antônio Gabriel Torres Cardoso<sup>1</sup> , José Marcos do Nascimento dos Santos Abreu<sup>1</sup>  &  
Reuber Albuquerque Brandão<sup>1</sup> 

The increased human population and alterations of the environment have made changes in land cover and have differential impacts on biodiversity (Hansen *et al.* 2013, Ordeñana *et al.* 2010). Some anthropogenic activities offer new vacant habitats that may be explored by generalist species (Howell *et al.* 2019). The preponderance of these opportunistic species over local fauna leads to biological homogenization (Vitule *et al.* 2021). The result are communities formed by supertramp species that can thrive under a wide array of conditions, as well as have high growth rates relative to the availability of the local resources (Clavel *et al.* 2011). Unfortunately, the ecological consequences of the biotic homogenization process may be significant and can lead to reductions in overall community and ecosystem function, stability, and adaptability (Olden *et al.* 2004). In the Distrito Federal, Brazil it is no different. According to the goal to develop and colonize the interior of Brazil as a public policy

during the '50s (Vidal 2009), the local vegetation cover and land use have been altered (UNESCO 2002), involving the replacement of native vegetation by pasture, agriculture, dams, urbanization, and highways (Ferreira 2010).

*Boana raniceps* (Cope, 1862) is a generalist medium-sized neotropical hylid characterized by a yellowish dorsum and by the presence of stripes on the thighs and inguinal regions (Faivovich *et al.* 2005). The species occurs from the north of the Amazon, passing through the Cerrado, Caatinga and reaching the Argentine and Paraguayan Chacos (Frost 2017) and is found in permanent or temporary water bodies associated with the gallery forest and natural and anthropized open environments (Vaz-Silva *et al.* 2020). *Boana raniceps* is the most ubiquitous and conspicuous hylid frog in the hotter and humid lowlands of the Cerrado landscape (Pacheco *et al.* 2018), emitting its load and

<sup>1</sup> Laboratório de Fauna e Unidades de Conservação, Departamento de Engenharia Florestal, Universidade de Brasília, 70910-900, Brasília, Distrito Federal, Brazil.

<sup>2</sup> Laboratório de Sistemática e Biogeografia de Anfíbios e Répteis – Manguari; Programa de Pós-Graduação em Biologia Animal, Instituto de Biociências, Universidade Federal de Mato Grosso do Sul, 79070-900, Campo Grande, Mato Grosso do Sul, Brazil.

<sup>3</sup> Programa de Pós-Graduação em Zoologia, Instituto de Biologia, Universidade de Brasília, 70910-900, Brasília, Distrito Federal, Brazil.

<sup>4</sup> Laboratório de Planejamento para Conservação da Biodiversidade; Programa de Pós-Graduação em Ecologia, Instituto de Ciências Biológicas, Universidade de Brasília, 70910-900, Brasília, Distrito Federal, Brazil.

\* Corresponding author: [beatrizdiogov@gmail.com](mailto:beatrizdiogov@gmail.com)

characteristic call usually during the rainy season (Vaz-Silva *et al.* 2020).

The herpetofauna of Distrito Federal (DF) is one of the most studied in the Cerrado biome (Nogueira *et al.*, 2010), but *B. raniceps* was never inventoried in any of the former herpetofauna works within the DF (Moreira & Barreto 1996, Brandão & Araujo 1998, Brandão & Araujo 2001, Brandão *et al.* 2006, Colli *et al.* 2011, Crema *et al.* 2014, Vaz-Silva *et al.* 2020). Herein, we report the first records of the species for the Distrito Federal and its possible recent colonization in the region.

We recorded *Boana raniceps* in three locations in Distrito Federal. We found two adult individuals (respectively CRC = 74 millimeters, 28

grams, and CRC = 87mm, 45,5g) during active night search on March 29, 2016 and December 26 of the same year in Lagoa Bonita, Estação Ecológica de Águas Emendadas, Planaltina-DF (15°35'15.80"S; 47°41'54.81"W, 959 meters above sea level). The site has open vegetation with the presence of buriti palms (*Mauritia flexuosa*), a predominance of invasive grasses and it has old, abandoned fish farms that are currently silted and colonized by several aquatic plants. It's worthwhile to cite that Lagoa Bonita is one of the most studied areas for the DF anurans. The individuals were found in buriti palms, where both were photographed (Figure 1), captured, measured, and released.



**Figure 1.** *Boana raniceps* individual near Lagoa Bonita, Estação Ecológica de Águas Emendadas, Planaltina-DF. Antonio G. T. Cardoso.

On another field trip, on December 19, 2019, we found a *Boana raniceps* individual in a bush near an artificial lake at Fazenda Mangabeiras, Planaltina-GO (15°28'54.15"S; 47°51'57.13"W, 741 m a.s.l.), integrated into the Integrated Economic Development Region of Distrito Federal. At the time, it was possible to hear more individuals vocalizing. The area where the species was found had many shrub and arboreal plants and is located in the farm Legal Reserve, a zone for rural properties intended for vegetation conservation and ecological tourism (Brasil 2012).

On May 18, 2021, the team of the Brasília Botanical Garden, Brasília-DF (15°52'32.4"S; 47°50'13.0"W, 1116 m a.s.l.), was managing the *Cyperus papyrus* L. stands in a small lake when they found another *Boana raniceps* individual, hidden in the dry foliage.

Despite being quite abundant in the state of Goiás (Vaz-Silva 2020), *Boana raniceps* had no records in the Distrito Federal until now. The species commonly colonizes anthropic habitats (Souza *et al.* 2010), using the available food sources and new habitats and breeding environments (Collins & Fahrig 2017). Possibly these records refer to new and recent colonization of the species in the region, since the areas where the species was found are in or near anthropized environments and the species is associated to permanent or temporary water bodies associated to gallery forest and natural and anthropized open environments (Vaz-Silva *et al.* 2020). We are unaware if the species was benefited by recent habitat changes in the Distrito Federal caused by the deep removal of natural habitats during the region colonization in the recent decades (Scariot *et al.* 2005). As a geographically widespread species, *B. raniceps* is subjected to a broad range of climatic conditions (Bonnetfond *et al.* 2020), so

another possibility is related to changes in local climate, caused by recent increments in Distrito Federal temperatures in the last decades (Hofmann *et al.* 2021).

#### ACKNOWLEDGMENTS

We thank the team that manages the Jardim de Cheiros, in the Jardim Botânico de Brasília, Brasília-DF, Millena Castro Ribeiro, Lucas de Albuquerque Miranda and Maria Rosa Vargas Zanatta for the partnership that complemented the writing of this scientific note. RAB thanks to Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for productivity grant (process #306644/2020-7). BDV and ERC thanks Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES) – Funding code 001.

#### REFERENCES

- Bonnetfond, A., Courtois, E.A., Sueur, J., Sugai, L.S.M. & Llusia, D. (2020). Climatic breadth of calling behavior in two widespread Neotropical frogs: Insights from humidity extremes. *Global Change Biology* 26:5431–5446.
- Brandão, R.A. & Araújo, A.F.B. (1998) A herpetofauna da Estação Ecológica de Águas Emendadas. In: Marinho-Filho, J., Rodrigues, F. & Guimarães, M. (Eds.) *Vertebrados da Estação Ecológica de Águas Emendadas: História Natural e Ecologia em um fragmento de cerrado do Brasil Central*. Instituto de Ecologia e Meio Ambiente do Distrito Federal, Brasília, pp 9–21.
- Brandão, R.A. & Araújo, A.F.B. (2001) A herpetofauna associada às matas de galeria no Distrito Federal. In: Ribeiro, J.F., Fonseca, C.E.L. & Sousa-Silva, J.C. (Eds.) *Cerrado: Caracterização e Recuperação de Matas de Galeria*. Empresa Brasileira de Pesquisa Agropecuária - Embrapa Cerrados & Ministério do Meio Ambiente, Planaltina, pp. 559–604.
- Brandão, R.A., Sebben, A. & Zerbini, G.J. (2006) A Herpetofauna da Área de Proteção Ambiental do Cafuringa. In: Secretaria de Meio Ambiente e Recursos Hídricos (Eds.) *A Área de Proteção Ambiental do*

- Cafuringa: A Última Fronteira Natural do Distrito Federal*. SEMARH, Brazil, pp 241–248.
- Brasil (2012) Lei nº 12.651, de 25 de maio de 2012. Presidência da República. Disponível em: [http://www.planalto.gov.br/ccivil\\_03/Ato2011-2014/2012/Lei/L12651.htm#art83](http://www.planalto.gov.br/ccivil_03/Ato2011-2014/2012/Lei/L12651.htm#art83) (acesso: 23 Setembro 2021).
- Colli, G., Nogueira, C.C., Leite, D.P., Ledo, R., Costa, B.M. & Brandão, R.A. (2011) Herpetofauna da Reserva Ecológica do IBGE e seu entorno. In: Ribeiro, M.L. (Ed.) *Reserva Ecológica do IBGE: Biodiversidade Terrestre*. IBGE Coordenação de Recursos Naturais e Estudos Ambientais, Rio de Janeiro, pp 132-145.
- Collins, S.J. & Fahrig, L. (2017) Responses of anurans to composition and configuration of agricultural landscapes. *Agriculture, Ecosystems & Environment* 239:399–409.
- Clavel, J., Julliard, R. & Devictor, V. (2011) Worldwide decline of specialist species: toward a global functional homogenization? *Frontiers in Ecology and the Environment* 9:222–228.
- Crema, A., de Souza, F.A.C.F., Patelli, L.F.P., de Carvalho, R.M.V. & Mesquita, D.O. (2014) Diversidade e distribuição de anfíbios anuros em matas de galeria do Distrito Federal, Brasil. *Revista Nordestina de Biologia* 23:3–27.
- Faivovich, J., Haddad, C.F.B., Garcia, P.C., Frost, D.R., Campbell, J.A. & Wheeler, W.C. (2005) Systematic review of the frog family Hylidae, with special reference to Hylinae: phylogenetic analysis and taxonomic revision. *Bulletin of the American Museum of Natural History* 17:1–240.
- Ferreira, I.C.B. (2010) Brasília: mitos e contradições na história de Brasília. In: Paviani, A., Jatobá, S.U., Cidade, L.C.F. & Barreto, F.F.P. (Eds.) *Brasília 50 anos: da capital a metrópole*. Universidade de Brasília, Brasília, pp. 23–54.
- Frost, D.R. (2021) *Amphibian Species of the World: an online reference*, version 6.1. Disponível em: <https://amphibiansoftheworld.amnh.org/Amphibia/A-nura/Hylidae/Cophomantinae/Boana/Boana-raniceps> (acesso: 02 Outubro 2021).
- Hansen, M. C., Potapov, P.V., Moore, R., Hancher, M., Turubanova, S.A., Tyukavina, A., Thau, D., Stehman, S.V., Goetz, S.J., Loveland, T.R., Kommareddy, A., Egorov, A., Chini, L., Justice, C.O. & Townshend, J.R.G. (2013) High-resolution global maps of 21st-century forest cover change. *Science* 342:850–853.
- Hofmann, G.S., Cardoso, M.F., Alves, R.J.V., Weber, E.J., Barbosa, A.A., Toledo, P.M., Pontual, F.B., Salles, L.O., Hasenack, H., Cordeiro, J.L.P., Aquino, F.E. & Oliveira, L.F.B. (2021) The Brazilian Cerrado is becoming hotter and drier. *Global Change Biology* 27:4060–4073
- Howell, H.J., Mothes, C.C., Clements, S.L., Catania, S.V., Rothermel, B.B. & Searcy, C.A. (2019) Amphibian responses to livestock use of wetlands: new empirical data and a global review. *Ecological Applications* 0:e01976 <https://doi:10.1002/eap.1976>.
- Moreira, G. & Barreto, L. (1996) Alimentação e variação sazonal na frequência de capturas de anuros em duas localidades do Brasil Central. *Revista Brasileira de Zoologia* 13:313–320.
- Nogueira, C., Colli, G.R., Costa, G. & Machado, R.B. (2010) Diversidade de répteis Squamata e evolução do conhecimento faunístico no Cerrado. In: Diniz, I.R., Filho, J.M., Machado, R.B. & Cavalcanti, R.B. (Eds.) *Cerrado: Conhecimento científico quantitativo como subsídio para ações de conservação*. Thesaurus, Brasília, pp. 331–372.
- Olden, J. & Poff, N.L. (2003) Toward a mechanistic understanding and prediction of biotic homogenization. *The American Naturalist* 162:442–460.
- Ordeñana, M.A., Crooks, K.R., Boydston, E.E., Fisher, R.N., Lyren, L.M., Siudyla, S., Haas, C.D., Harris, S., Hathaway, S.A., Turschak, G.M., Miles, A.K. & Van Vuren, D.H. (2010) Effects of urbanization on carnivore species distribution and richness. *Journal of Mammalogy* 91:1322–1331.
- Pacheco, E.O., Mângia, S. & Santana, D.J. (2018) Diversity and distribution of anurans among different vegetation physiognomies in a savannah landscape in Central Brazil. *Herpetology Notes* 11:255–262.

- Paula, A.D. (2012) *Estrutura e dinâmica de uma comunidade de anuros no hotspot de biodiversidade do Cerrado*. Universidade de Brasília, Brasília, 79pp.
- Santos, F.L., Uetanabaro, M., Landgref-Filho, P., Piatti, L. & Prado, C.P.A. (2010) Herpetofauna, municipality of Porto Murtinho, Chaco region, State of Mato Grosso do Sul, Brazil. *CheckList* 6:470–475.
- Scariot, A., Sousa-Silva, J.C. & Felfili, J.M. (2005) *Ecologia, Biodiversidade e Conservação*. Ministério do Meio Ambiente, Brasília, 118pp.
- UNESCO. (2002) *Vegetação do Distrito Federal: tempo e espaço*. UNESCO, Brasília, 92pp.
- Vaz-Silva, W., Maciel, N.M., Nomura, F., Morais, A.R., Batista, V.G., Santos, D.L., Andrade, S.P., Oliveira, A.A.B., Brandão, R.A. & Bastos, R.P. (2020) *Guia de identificação das espécies de anfíbios (Anura e Gymnophiona) do estado de Goiás e do Distrito Federal, Brasil Central*. Sociedade Brasileira de Zoologia, Curitiba, 223pp.
- Vidal, L. (2009) *De Nova Lisboa a Brasília: a invenção de uma capital (séculos XIX-XX)*. Universidade de Brasília, Brasília, 352pp.
- Vitule, J. R.S., Occhi, T.V.T., Carneiro, L., Daga, V.S., Frehse, F.A., Bezerra, L.A.V., Forneck, S., Pereira, H.S., Freitas, M.O., Hegel, C.G.Z., Abilhoa, V., Grombone-Guaratini, M.T., Queiroz-Sousa, J., Pivello, V.R., Silva-Matos, D.M., Oliveira, I., Toledo, L.F., Vallejos, M.A.V., Zenni, R.D., Ford, A.G.P. & Braga, R.R. (2021) Non-native Species Introductions, Invasions, and Biotic Homogenization in the Atlantic Forest. In: Marques, M.C.M. & Grelle, C.E.V. (Eds.) *The Atlantic Forest: History, Biodiversity, Threats and Opportunities of the Mega-diverse Forest*. Springer International Publishing, pp. 269–295.

Recebido em 15/10/2021

Aceito em 21/02/2022

Publicado em 24/02/2022



This is an open-access article distributed under the terms of the Creative Commons Attribution License.